

DEHN FILLINGS CREATING ESSENTIAL SPHERES AND TORI

SANGYOP LEE, SEUNGSANG OH, AND MASAKAZU TERAGAITO

ABSTRACT. Let M be a simple 3-manifold with a toral boundary component. It is known that if two Dehn fillings on M along the boundary produce a reducible manifold and a toroidal manifold, then the distance between the filling slopes is at most three. This paper gives a remarkably short proof of this result.

Let M be a *simple* 3-manifold with a toral boundary component $\partial_0 M$, that is, it contains no essential sphere, disk, torus or annulus. The *slope* of an essential, unoriented, simple closed curve on $\partial_0 M$ is its isotopy class. We assume that α and β are two slopes on $\partial_0 M$ such that $M(\alpha)$ is a reducible manifold and $M(\beta)$ contains an essential torus. The goal is to measure the upper bound for $\Delta(\alpha, \beta)$ (the minimal geometric intersection number of α and β), and Oh [Oh1] and Wu [Wu] independently gave the optimum upper bound 3 for this case. In the present paper we give another short proof of this result.

Theorem 1. *Let M be a simple 3-manifold such that $M(\alpha)$ is reducible and $M(\beta)$ is toroidal. Then $\Delta(\alpha, \beta) \leq 3$.*

Assume $\Delta(\alpha, \beta) \geq 4$ for contradiction. Let $\widehat{Q}$ be a reducing sphere in $M(\alpha)$ which intersects the filling solid torus V_α in a family of meridian disks. We choose $\widehat{Q}$ so that $q = |\widehat{Q} \cap V_\alpha|$ is minimal (over all reducing spheres $\widehat{Q}$ in $M(\alpha)$). Similarly let $\widehat{T}$ be an essential torus in $M(\beta)$ meeting in $t = |\widehat{T} \cap V_\beta|$ meridian disks, the number of which is minimal over all such tori. Let $Q = \widehat{Q} \cap M$ and $T = \widehat{T} \cap M$. By an isotopy of Q , we may assume that Q and T intersect transversely, and $Q \cap T$ has the minimal number of components. Then as described in [Oh1], we obtain graphs G_Q in $\widehat{Q}$ and G_T in $\widehat{T}$. We use the definitions and terminology of [Oh1, Section2].

Lemma 2. [BZ, Lemmas 2.3, 4.1, 4.3] $q \geq 3$ and $t \geq 3$.

Lemma 3. [GL2, Theorem 2.4] G_T cannot contain Scharlemann cycles on distinct label pairs.

Lemma 4. [Wu, Lemmas 1.4, 1.8] G_Q cannot have more than $\frac{t}{2} + 2$ mutually parallel edges connecting parallel vertices. Furthermore, if G_Q has $\frac{t}{2} + 2$ mutually parallel edges connecting parallel vertices, then $t \equiv 0 \pmod{4}$.

Theorem 5. [JLOT, Theorem 1.1] *If $M(\alpha)$ and $M(\beta)$ contain a projective plane and an essential torus respectively, then either $\Delta(\alpha, \beta) \leq 2$ or $\Delta(\alpha, \beta) = 3$ with $t = 2$.*

Now we will prove Theorem 1.

1991 *Mathematics Subject Classification.* 57M50.

Key words and phrases. Dehn filling, reducible manifold, toroidal manifold.

Proof. We distinguish two cases;

Case I) Suppose that there is a vertex x of G_Q such that more than $\frac{3}{2}t$ edges connect x to antiparallel vertices.

This implies that there exist more than $\frac{3}{2}t$ x -edges connecting parallel vertices in G_T . Consider the subgraph of G_T consisting all vertices and those all x -edges of G_T . An Euler characteristic count gives that it contains more than $\frac{1}{2}t$ disk faces whose boundaries are x -edge cycles of G_T . By [HM, Proposition 5.1], each disk face contains a Scharlemann cycle. By Lemma 3, these all Scharlemann cycles are, say, 12-Scharlemann cycles.

Construct a graph Γ in $\widehat{T}$ as follows. Choose a *dual vertex* in the interior of the face G_T bounded by a 12-Scharlemann cycle, and let the vertices of Γ be the vertices of G_T together with these dual vertices. The edges of Γ are defined by joining each dual vertex to the vertices of the corresponding Scharlemann cycle in the obvious way. Let s be the number of 12-Scharlemann cycles. Then $s > \frac{1}{2}t$. Γ has $t + s$ vertices and at least $3s$ edges because each Scharlemann cycle has order at least 3 by Theorem 5. Note that if G_T contains an Scharlemann cycle of length two, then $M(\alpha)$ contains a projective plane [GLi]. Again an Euler characteristic count guarantees that Γ has a disk face E . But E determines a 1-edge cycle bounding a disk face in E which, as long as $q > 2$, contains a Scharlemann cycle. This contradicts the definition of Γ .

Case II) As the negation of case I), suppose that each vertex x of G_Q has at least $(\Delta - \frac{3}{2})t$ labels where edges connecting parallel vertices are incident.

Let G_Q^+ be the subgraph of G_Q consisting of all vertices and edges connecting parallel vertices of G_Q . By the assumption, every vertex of G_Q^+ has valency at least $(\Delta - \frac{3}{2})t$. Let $\overline{G}_Q^+$ be the reduced graph of G_Q^+ . By Lemma 4, any vertex has valency at least 3 in $\overline{G}_Q^+$. So, we can choose an innermost component H of $\overline{G}_Q^+$, and a block $\overline{\Lambda}$ of H with at most one cut vertex of H . Let Λ be the subgraph of G_Q^+ corresponding to $\overline{\Lambda}$.

Note that there is a disk D in $\widehat{Q}$ such that $D \cap G_Q^+ = \Lambda$. A vertex of $\overline{\Lambda}$ is a *boundary vertex* if there is an arc connecting it to ∂D whose interior is disjoint from $\overline{\Lambda}$, and an *interior vertex* otherwise. Remark that $\overline{\Lambda}$ has an interior vertex by [Wu, Lemma 3.2].

If Λ contains a cut vertex of H , then let x be a label such that the cut vertex has an edge attached with label x there. Otherwise, let x be any label. Then each vertex of Λ , except a cut vertex, has at least two edges attached with label x . Thus we can find a great x -cycle. This implies that Λ has a Scharlemann cycle and so t is even by [BZ, Lemma 2.2(1)].

Let v, e and f be the numbers of vertices, edges and faces of its reduced graph $\overline{\Lambda}$. Also say v_i, v_∂ and v_c the numbers of interior vertices, boundary vertices and a cut vertex of $\overline{\Lambda}$. Hence $v = v_i + v_\partial$ and $v_c = 0$ or 1 . Since each face of $\overline{\Lambda}$ is a disk with at least 3 sides, we have $2e \geq 3f + v_\partial$. Combined with $1 = \chi(\text{disk}) = v - e + f$, we get $e \leq 3v - v_\partial - 3 = 3v_i + 2v_\partial - 3$.

Suppose that every interior vertex of $\overline{\Lambda}$ has valency at least 6 and that every boundary vertex except a cut vertex has valency at least 4. Since a cut vertex has valency at least 2, we have $2e \geq 6v_i + 4(v_\partial - v_c) + 2v_c$. These two inequalities give us that $0 \leq v_c - 3$, a contradiction.

Hence there are two cases. For the first case, assume that some boundary vertex y of $\overline{\Lambda}$ has valency at most 3. There are at least $(\Delta - \frac{3}{2})t$ edges in G_Q which are incident to y and connect y to parallel vertices. By Lemma 4, $\frac{5}{2}t \leq (\Delta - \frac{3}{2})t \leq 3(\frac{t}{2} + 2)$, i.e. $t \leq 6$ and $t \neq 6$.

But in the case of $t = 4$ there are at least 4 mutually parallel edges in Λ , which implies that G_Q contains two S -cycles with disjoint label pairs. Then $M(\beta)$ contains a Klein bottle as in the proof of [GL1, Lemma 3.10]. (Note that $M(\beta)$ is irreducible [GL2]. Hence the edges of an S -cycle cannot lie in a disk on $\widehat{T}$.) This contradicts to [Oh2, Theorem 1.1].

For the second case, assume that some interior vertex of $\bar{\Lambda}$ has valency at most 5. Again we have $4t \leq \Delta t \leq 5(\frac{t}{2} + 2)$. Then we can use the same argument as above. $\square$

REFERENCES

- [BZ] S. Boyer and X. Zhang, *Reducing Dehn filling and toroidal Dehn filling*, Topology Appl. **68** (1996), 285–303.
- [CGLS] M. Culler, C. Gordon, J. Luecke and P. Shalen, *Dehn surgery on knots*, Ann. Math. **125** (1987), 237–300.
- [GL1] C. Gordon and J. Luecke, *Dehn surgeries on knots creating essential tori, I*, Communications in Analysis and Geometry **3** (1995), 597–644.
- [GL2] C. Gordon and J. Luecke, *Reducible manifolds and Dehn surgery*, Topology **35** (1996), 385–409.
- [GLi] C. Gordon and R. Litherland, *Incompressible planar surfaces in 3-manifolds*, Topology Appl. **18** (1984), 121–144.
- [HM] C. Hayashi and K. Motegi, *Only single twists on unknots can produce composite knots*, Trans. Amer. Math. Soc. **349** (1997), 4465–4479.
- [JLOT] G. T. Jin, S. Lee, S. Oh and M. Teragaito, *P^2 -reducing and toroidal Dehn fillings*, preprint.
- [Oh1] S. Oh, *Reducible and toroidal 3-manifolds obtained by Dehn fillings*, Topology Appl. **75** (1997), 93–104.
- [Oh2] S. Oh, *Dehn filling, reducible 3-manifolds, and Klein bottles*, Proc. Amer. Math. Soc. **126-1** (1998), 289–296.
- [Wu] Y.Q. Wu, *Dehn fillings producing reducible manifolds and toroidal manifolds*, Topology **37** (1998), 95–108.

DEPARTMENT OF MATHEMATICS, KOREA ADVANCED INSTITUTE OF SCIENCE AND TECHNOLOGY, TAEJON 305–701, KOREA

E-mail address: apple@mathx.kaist.ac.kr

DEPARTMENT OF MATHEMATICS, CHONBUK NATIONAL UNIVERSITY, CHONJU, CHONBUK 561-756, KOREA

E-mail address: soh@math.chonbuk.ac.kr

DEPARTMENT OF MATHEMATICS AND MATHEMATICS EDUCATION, HIROSHIMA UNIVERSITY, KAGAMIYAMA 1-1-1, HIGASHI-HIROSHIMA 739-8524, JAPAN

E-mail address: teragai@hiroshima-u.ac.jp